

Mobility Status Following Subtotal Calcaneotomy in Diabetic Foot Related Hindfoot Infection

Huang S^{1*}, Shi CJY¹, Seng DWR², Premchand AXR¹ and Park DH¹

¹Department of Orthopaedic Surgery, Khoo Teck Puat Hospital, Singapore

²Department of Orthopaedic Surgery, Woodlands Health Campus, Singapore

*Corresponding author:

Shirong Huang,
Department of Orthopaedic Surgery, Khoo Teck
Puat Hospital, Singapore

Received: 06 Apr 2024

Accepted: 20 May 2024

Published: 25 May 2024

J Short Name: ACMCR

Copyright:

©2024 Huang S. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially

Keywords:

Diabetic foot ulcer; Calcaneotomy; Ambulation;
Limb preservation

Citation:

Huang S, Mobility Status Following Subtotal Calcaneotomy in Diabetic Foot Related Hindfoot Infection. Ann Clin Med Case Rep. 2024; V13(17): 1-5

1. Abstract

Diabetic foot related infections are the main causes of major amputations in diabetic patients [1]. In particular, hindfoot ulcers and infections are often harder to treat. A subtotal calcaneotomy can be a viable alternative to major amputation in diabetic related hindfoot infections. We conducted a retrospective study of a cohort of diabetic patients undergoing a subtotal calcaneotomy in our institution. All patients underwent an open subtotal calcaneotomy for diabetic foot related hindfoot infections. From January 2011 to October 2021, a total of 13 patients were identified. The mean follow-up period was 20 [2] months. Of the 13 patient analysed, 5 maintained their mobility status post subtotal calcaneotomy.

A subtotal calcaneotomy is a viable limb salvage alternative to major lower limb amputations in diabetic related hindfoot infections, with a significant proportion of patients maintaining their pre-operative mobility statuses, which is comparable to currently published data on major lower limb amputations.

2. Introduction

Patients with diabetic foot related ulcers in the hindfoot are at particular risk of developing recalcitrant heel ulcers progressing to calcaneal osteomyelitis. The major pathologies of neuropathy, ischaemia, and infection often co-exist and exacerbate the condition. Healing of such hindfoot ulcers is also impaired by the limited amount of soft tissue coverage over the calcaneus. Major amputation rates and mortality of such hindfoot infections are higher compared to those in the forefoot or midfoot [2], with a study re-

porting a 52% rate of major amputation in heel osteomyelitis [3]. The prevalence of diabetes in the Singaporean adult population is estimated to be at 8.6% in 2017 [4]. Diabetic related complications, along their subsequent economical and healthcare associated costs, have been well documented [5,6]. Diabetic related foot infections can be a significant source of these costs as they often result in major lower limb amputation [7]. Locally in Singapore, diabetic related major lower limb amputation has been reported to be at a rate of 95 per 10,000 [8]. Major lower limb amputations have been proven to result in significant morbidity and mortality [9,10]. Of patients undergoing a below knee amputation, half are ambulant at 1 year post-amputation [9,11].

It has been established that a calcaneotomy is a viable alternative to major amputation in diabetic related hindfoot infections, with preservation of ambulatory status in a higher proportion of patients [12-14,17,18] compared to patients undergoing major lower limb amputations [9,11,15]. Minor complication rates following a calcaneotomy have been quoted at 24%, with approximately 10% of patients ultimately requiring a major lower limb amputation [14].

The aim of our study is to present the ambulatory outcomes of patients undergoing a subtotal calcaneotomy for diabetic hindfoot infections.

3. Methods

Our study received approval from the Ethics Review Board of our Institution. We performed a retrospective review of the case notes of all patients admitted to our hospital from January 2012 to Oc-

tober 2021 who presented with diabetic hindfoot infections and underwent subtotal calcaneotomies. All authors had no conflicts of interest to declare.

The inclusion criteria the study were diabetic patients with clinical evidence of hindfoot soft tissue infection, with or without clinical or radiological evidence of calcaneal osteomyelitis. They also were pre-morbidly ambulant with or without the use of walking aids.

Patients who met the following exclusion criteria were removed from the study population; patients who underwent a conversion to major lower limb amputation, or those who had follow-up data of less than 6 months post-operatively. Patients with significant comorbidities like end stage renal failure or peripheral arterial disease were not excluded, even though these patients were at a higher risk of wound healing complications.

The patient parameters that were collected included demographics (age, gender, ethnicity), medical comorbidities, pre-morbid and post-operative mobility statuses, time to wound healing, as well as details of their surgical procedures.

All details were obtained via screening of electronic medical records. Their pre-operative and post-operative ambulatory status, as assessed by a combination of allied health professionals (physiotherapists, occupational therapists) in our institution, was also compared. The assessment by the allied health professionals were performed both during their inpatient stay, as well as during outpatient therapy follow-ups. The assessment of each patient's ambulation status was collected at a minimum duration of 6 months post-operatively. Our primary outcome measures were ambulatory status post-operatively, and time to complete wound healing.

All calcaneotomies were done via an open approach, usually incorporating any open wounds present. Definitive calcaneotomies were usually performed after multiple debridements to remove dead and infected tissue to achieve initial source control. Patients were operated on under a general or regional anaesthetic, with no tourniquet control and were either positioned prone or lateral decubitus. All infected and non-viable tissue were removed. The amount of calcaneus resected depended on clinical evaluation of the soft tissue condition, or was based off the extent of calcaneal osteomyelitis as evaluated on advanced radiological imaging (MRI).

The primary outcome measure used was return to mobility as determined by reports from clinical review or from professional allied health assessments.

2 patients had a supplementary external circular frame applied after their subtotal calcaneotomies. 3 patients underwent an antero-lateral thigh (ALT) flap following their calcaneotomies.

4. Results

A total of 28 patients underwent subtotal calcaneotomies for diabetic related hindfoot infections in our hospital from January 2011 to October 2021. 15 patients were excluded from the study due to conversion to below knee amputation (6 patients), inadequate duration of follow-up (7 patients), and pre-morbidly non-ambulant patients (2 patients). A total of 13 patients were included in the final analysis. Patients were followed up for a minimum of 6 months post-operatively. The duration of follow-up ranged from 6 to 96 months, with a mean follow-up duration of 20.2 months. Table 1 shows the characteristics of the study population.

We classified the ambulatory statuses for the study population into 2 broad categories; ambulant and non-ambulant. Within the ambulant population, this was further sub-divided into 3 groups; those ambulant without a walking aid, those requiring a walking stick, and those requiring a walking frame. The non-ambulant group comprised of patients who were pre-operatively wheelchair or bedbound.

Of the 13 patients analysed, 5 (38.5%) maintained their pre-operative mobility after a subtotal calcaneotomy. There were no patients who experienced an improvement in their ambulatory status. Notably, 2 of the 8 patients whose ambulatory status declined post-operatively also had subsequent contralateral below knee amputations which significantly affected mobility. Another had an ipsilateral transmetatarsal amputation performed for a midfoot infection. Hence, the proportion of patients maintaining their ambulatory status with an isolated subtotal calcaneotomy was 5 out of 10 patients (50%).

Of the 13 patients, 9 (69.2%) had documented full wound healing. The mean time for full wound healing was 8.33 months. Of the 4 patients whose wounds did not heal, 2 had undergone an ALT flap procedure, with documented flap complications at the time of data collection. 1 patient demised at 11 months postop, prior to documented wound healing.

6 out of 28 patients (21.4%) required conversion to a below knee amputation following their calcaneotomy. Out of the 5 patients who maintained their ambulatory statuses, only 2 (40%) had documented peripheral vascular disease. In addition, only 1 out of 5 (20%) had end stage renal failure. Table 2 showcases the overall pre and post-operative ambulatory status for our study population. Table 3 showcases the breakdown of the change in ambulatory statuses for each category.

Table 1: Characteristics of Study Population (patients undergoing subtotal calcaneotomy following diabetic hindfoot related infection)

Characteristics	Total Study Population (n=13)
Male	8
Female	5
Age (years), mean	57.2
Ethnicity	
- Chinese	9
- Malay	2
- Indian	2
- Others	0
Ethnicity (% breakdown)	
- Chinese	69.2
- Malay	15.4
- Indian	15.4
Co-morbidities	
Peripheral vascular disease	8
End stage renal failure	5
Ischaemic heart disease	5
Stroke	2

Table 2: Pre-operative and post-operative ambulatory statuses

Pre-operative ambulatory status	n=13	Post-operative ambulatory status	n=13
Ambulant	13	Ambulant	7
- Without aid	8	- Without aid	4
- With walking stick	1	- With walking stick	2
- With walking frame	4	- With walking frame	1
Non-ambulant	0	Non-ambulant	6

Table 3: Change in ambulatory status post subtotal calcaneotomy

Maintained (n=5)	Ambulant without aid	4
	Walking stick	1
	Walking frame	0
Decreased (n=8) Ø = Post-operative	Premorbid: Ambulant without aid (n=4)	
	Ø Walking stick	1
	Ø Walking frame	1
	Ø Non-ambulant	2
	Premorbid: Ambulant with walking stick (n=0)	
	Ø Walking frame	0
	Ø Non-ambulant	0
	Premorbid: Ambulant with walking frame (n=4)	
	Ø Non-ambulant	4

5. Discussion

Our study population is fairly representative of the overall Singaporean population in terms of the ethnic distribution. The 2020 Singapore Population Census is as such; the Singapore ethnic breakdown is 74.3% Chinese, 13.5% Malay, 9% Indian, with 3.2% classified as others. This is comparable to our study population distribution, with a slight increased representation in the Indian racial group.

There have been limited studies on the ambulatory outcomes following a calcaneotomy for hindfoot related infections. These studies suggest that ambulatory outcomes following a calcaneotomy are more favourable compared to currently published data on below knee amputations. In particular, a systematic review by Shade in 2012 revealed that 85% of post-calcaneotomy patients either maintained or improved their ambulatory statuses post-operatively.¹⁴ While our study results do not showcase such a similarly high proportion of patients maintaining their ambulatory statuses, a 50% proportion is comparable to currently published literature on post below knee amputees and their ambulatory statuses post-operatively with a prosthesis.

The healing rates showcased in our study is also comparable to locally published data on below knee amputations. In a prospective cohort study by Wong in 2013, 73.5% of below knee amputees had good stump healing within 6 months, and did not require more proximal amputations.⁹ In comparison, 9 out of our 13 patients (69.2%) had documented full wound healing, with a mean healing time of 8.33 months. This further demonstrates that the subtotal calcaneotomy has comparable wound healing rates compared to a below knee amputation, and is a viable limb salvage alternative to major lower limb amputations.

Our study also had a good representative cohort of patients, with good follow-up duration. We included patients with significant medical co-morbidities, and a spectrum of pre-operative ambulatory statuses. Patients with complex medical co-morbidities such as those with peripheral vascular disease or end stage renal failure were not excluded. This is highly reflective of the actual demographic of patients presenting with diabetic hindfoot related infections. The mean follow-up duration of 20.2 months also provided us with adequate data when it came to post-operative ambulatory as well as wound assessments.

However, our study is also limited due to the heterogeneity of our study population. As a retrospective study, we were unable to effectively control factors such as co-morbidities, wound size, surgical technique, and post-operative protocols such as antibiotic therapy, offloading footwear/insoles, or weightbearing status. For

example, the presence of peripheral vascular disease or end stage renal failure could negatively impact wound healing and therefore ambulatory status post-operatively. Poor pulses or a low ankle brachial pressure index have been shown to be associated with poor clinical outcomes following below knee amputations.⁹ The lack of a standardised protocol following the subtotal calcaneotomies in our series could have a potential impact on the interpretation of our results, as the escalation of the weightbearing status over the ipsilateral limb post-operatively would vary from case to case. This would undoubtedly affect the progress of physical therapy post-operatively, and therefore the potential recovery of ambulation.

The low numbers in our study is also a limiting factor. Patients with diabetic related hindfoot infections are a difficult subset amongst those with diabetic foot infections, thus our institution saw a low volume of patients undergoing a subtotal calcaneotomy who fit our inclusion and exclusion criteria. The necessary exclusion criteria significantly reduced the number of patients eligible for analysis. A significant number of patients, 6 out of 28 (21.4%), were excluded due to conversion to below knee amputation, thus limiting the utility of evaluating their post-operative mobility for the purpose of this study. 2 out of 28 (7.1%), were also non-ambulant pre-operatively and were hence excluded from analysis.

A number of our patients also underwent subsequent coverage procedures, which included ALT flaps. Out of the 3 patients in our study who underwent ALT flap coverage procedures, only 1 had documented healing at 16 months post-calcaneotomy. We propose that these additional coverage procedures may have delayed the weightbearing statuses for these patients, and may have made post-operative mobilisation more challenging. In addition, these coverage procedures are likely also associated with patients who have had a more extensive infection, and therefore a larger post-operative soft tissue defect.

6. Conclusion

A subtotal calcaneotomy is a viable limb salvage alternative to major lower limb amputations in diabetic related hindfoot infections, with a significant proportion of patients (50 %) maintaining their pre-operative mobility statuses compared to currently published data on major lower limb amputations. Nearly 70% of our patients also had documented full wound healing, with a mean healing time of 8.33 months. Our case series also showed that approximately 21% of patients undergoing a subtotal calcaneotomy eventually required a below knee amputation. Further studies may be needed on the rates of morbidity and mortality following a calcaneotomy to further evaluate its role as an alternative to major lower limb amputations.

References

1. Pecoraro RE, Reiber GE, Burgess EM. Pathways to diabetic limb amputation: basis for prevention. *Diabetes Care*. 1990; 13: 513-21.
2. Pickwell KM, Siersma VD, Kars M, Holstein PE, Schaper NC, Eurodiale consortium. Diabetic foot disease: impact of ulcer location on ulcer healing. *Diabetes Metab Res Rev*. 2013; 29(5): 377-83.
3. Faglia E, Clerici G, Caminiti M, Curci V, Somalvico F. Influence of osteomyelitis location in the foot of diabetic patients with transtibial amputation. *Foot Ankle Int*. 2013; 34(2): 222-7.
4. Ministry of Health, Singapore. Executive summary on national population health survey 2016/17, 2017.
5. Schirr-Bonnans S, Costa N, Derumeaux-Burel H, Bos J, Lepage B, Garnault V, et al. Cost of diabetic eye, renal and foot complications: a methodological review. *Eur J Health Econ*. 2017; 18(3): 293-312.
6. WHO. Global report on diabetes. 2016.
7. Pecoraro RE, Reiber GE, Burgess EM. Pathways to diabetic limb amputation: basis for prevention. *Diabetes Care*. 1990; 13: 513-21.
8. Riandini T, Pang D, Toh MPHS, Tan CS, Choong AMTL, Lo ZJ, et al. National Rates of Lower Extremity Amputation in People With and Without Diabetes in a Multi-Ethnic Asian Population: a Ten Year Study in Singapore. *Eur J Vasc Endovasc Surg*. 2022; 63(1): 147-55.
9. Wong KL, Nather A, Liang S, Chang Z, Wong TTC, Lim CT. Clinical Outcomes of Below Knee Amputations in Diabetic Foot Patients. *Ann Acad Med Singapore* 2013; 42: 388-94.
10. Aulivola B, Hile CN, Hamdan AD, Sheahan MG, Veraldi JR, Skillman JJ, et al. Major Lower Extremity Amputation: Outcome of a Modern Series. *Arch Surg*. 2004; 139(4): 395-9.
11. Chopra A, Azarbal AF, Jung E, Abraham CZ, Liem TK, Landry GJ, et al. Ambulation and functional outcome after major lower extremity amputation. *Journal of Vascular Surgery* 2018; 67(5): 1521-29.
12. Smith DG, Stuck RM, Ketner L, Sage RM, Pinzur MS. Partial calcaneotomy for the treatment of large ulcerations of the heel and calcaneal osteomyelitis. An amputation of the back of the foot. *J Bone Joint Surg Am*. 1992; 74(4): 571-6.
13. Bollinger M, Thordarson DB. Partial calcaneotomy: an alternative to below knee amputation. *Foot Ankle Int*. 2002; 23(10): 927-32.
14. Schade VL. Partial or Total Calcaneotomy as an Alternative to Below-the-Knee Amputation for Limb Salvage: A Systematic Review. *J Am Podiatr Med Assoc*. 2012; 102(5): 396-405.
15. Sansosti LE, Crowell A, Choi ET, Meyr AJ. Rate of and Factors Associated with Ambulation After Unilateral Major Lower-Limb Amputation at an Urban US Tertiary-Care Hospital with a Multidisciplinary Limb Salvage Team. *J Am Podiatr Med Assoc*. 2017; 107(5): 355-64.
16. Dalla Paola L, Carone A, Boscarino G, Scavone G, Vasilache L. Combination of Open Subtotal Calcaneotomy and Stabilization With External Fixation as Limb Salvage Procedure in Hindfoot-Infected Diabetic Foot Ulcers. *Int J Low Extrem Wounds*. 2016; 15(4): 332-7.
17. Baravarian B, Menendez MM, Weinheimer D, Lowery C, Kosanovich R, Vidt L. Subtotal calcaneotomy for the treatment of large heel ulceration and calcaneal osteomyelitis in the diabetic patient. *The Journal of foot and ankle surgery: official publication of the American College of Foot and Ankle Surgeons*. 1999; 38(3): 194-202.
18. Oliver NG, Steinberg J, Powers KA, Evans KK, Kim PJ, Attinger CE. Lower Extremity Function following Partial Calcaneotomy in High-Risk Limb Salvage Patients. *Journal of Diabetes Research*. 2015; 2015: 432164.