

The Mooji Effect of “Osborne-Cotterill Lesion”. New Approach to Old Concept

Mojahed (Mooji) Sakhnini MD, MPH, EBHS, FRCSEd, FEBOT

Hand and Microsurgery Specialist, Israel

*Corresponding author:

Sakhnini Mojahed,
Hand and Microsurgery Specialist, Israel
E-mail: Mojahed.sakh@gmail.com

Received: 26 Oct 2024

Accepted: 13 Nov 2024

Published: 19 Nov 2024

J Short Name: ACMCR

Copyright:

©2024 Sakhnini Mojahed. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially

Keyword:

“Osborne-Cotterill Lesion”; Elbow dislocation;
Reconstruction; Ligament; Stability

Citation:

Sakhnini Mojahed, The Mooji Effect of “Osborne-Cotterill Lesion”. New Approach to Old Concept. Ann Clin Med Case Rep. 2024; V14(10): 1-8

1. Abstract

1.1. Introduction

“Osborne-Cotterill Lesion” leaves an impression in the capitellum following injury or dislocation of the elbow. The clinical importance of this lesion is not yet recognized. We suggest a simple algorithm to measure its importance clinically and offer surgical treatment with bone graft and plating to prevent engagement of the radial head in this lesion calling this process of engagement “Mooji effect”.

1.2. Methods

we present two cases that demonstrates the surgical strategy for approaching “Osborne-Cotterill Lesion” when potential engagement is imminent. We also assign critical point in the capitellum that if breached then our surgical plan is employed.

1.3. Results

For one case, excellent result was achieved, and the other case was lost to follow-up. When a critical point is used in the context of “Osborne-Cotterill Lesion” as indicator for instability, then straightforward decision could be made when to proceed with our suggested strategy – bone grafting and plating “Osborne-Cotterill Lesion”.

1.4. Conclusion

We recommend specific surgical treatment for “Osborne-Cotterill Lesion” when indicated following our algorithm to prevent the “Mooji effect” and its repercussions including instability and limited range of motion.

2. Introduction

Elbow dislocation is a common injury that in majority of cases

is treated conservatively with good outcome [1,2]. Nevertheless, there are certain characteristics that should be considered to optimize outcome and prevent complications, mainly instability and pain [3]. Some of the risk factors are not under control like laxity, age and some are related to sport activities [4]. But what concerns us is the injury character, meaning if there is an impaction injury to the capitellum or fracture of the anterior lip of the radius that otherwise would be treated conservatively. We promote further discerning of these patterns and analyzing it to prevent residual instability that could eventually affect quality of life. In the literature “Osborne-Cotterill Lesion” is defined as a depression fracture in the capitellum [5,6]. Subsequent literature tried to address this injury and relate the depression in the capitellum to instability and described the depression as an injury caused by radial head dislocation impinging upon the capitellum and causing a crater [7]. Furthermore, a question was asked repeatedly about the importance of this lesion regarding instability raising the question whether part of the instability is related to this osseous defect [8]. Some authors tried to answer this question with anecdotal cases, case series or biomechanically in laboratory conditions [9-12]. What is interesting is that description of the fragment is nearly always posterior and lateral and not central in the capitellum which indicates that the fragment is produced when the radial head dislocates laterally and posteriorly and hits the capitellum at this specific location in a predicted manner with the same mechanism [6,7]. The lesion could cause engagement of the radial head, and the radial head may be dislocated and stuck in that position necessitating open reduction [13]. As the process of dislocation could lead to “Osborne-Cotterill Lesion” with impaction injury, we outline the engagement process of the radial head when it slips in that lesion, calling it – “The

Mooji effect”. We assign importance to this effect clinically and biomechanically and suggest addressing it in a tailored and specific manner. Otherwise, even with surgical intervention that intends to reanimate LUCL with re-insertion or reconstruction, the “Mooji effect” will lead to instability translated into dislocation or limited range of motion to prevent engagement [11]. In this series we represent a different approach to treat this enigmatic problem. We suggest bone grafting the “Osborne-Cotterill Lesion” and adding a posterior plate to lock the graft and to further thicken the barrier for dislocation. We will discuss this strategy with the help of two cases and present a simple algorithm to help decide when to address “Osborne-Cotterill Lesion” surgically.

3. The Critical Point for “Mooji Effect”

The anatomy of the radio-capitellar joint is intriguing; There is a match between the radius of radial head and the radius of the capitellum as this joint perform a 3D motion- Flexion-extension, pro-supination and translation[14]. Extension could be accompanied by supination movement and translation at the same time. For the radial head to dislocate, it must pass a critical point at the posterior edge of the capitellum. Passing that point it could relocate back or engage a shallow depression in the capitellum. To decide where this critical point is, we make several hypothetical points:

1. The diameter of the capitellum is the same diameter of the radial head [15].
2. The dislocation includes a translation of the radial head and not only a pendulum movement of the radial head against the capitellum [16].
3. The capitellum functional surface is inclined in average 45 degrees relative to the humerus [17].

4. The hyaline cartilage of the capitellum-functioning surface-covers 180 degrees (half of a full sphere).
5. The radial head has intrinsic translation movement of 1 radian up to the bisecting point of the capitellum when the posterior capsule is injured (following fracture or dislocation) [12,18].

These assumptions could be partly referred from literature and partly from observation and mechanical deduction. Based on these assumptions, we build the safe zone that indicates the distance the radial head could travel before it passes the critical point.

In full extension the posterior edge of the radial head passes this critical point (Figure 1). This is because the capitellum being a half sphere is anteriorly sloped. Because of the slope of the capitellum, the critical point forms a 45-degree angle between the center of the capitellum and a vertical line parallel to the humerus or horizontal line parallel to the radial head in neutral full extension. We define “ α ” angle as 2 radians (114.6 degrees) of the capitellar sphere equal to diameter of the radial head which is the distance the anterior edge of the radius needs to pass before dislocating when it is in full extension. Form “ α ” we must take out “ β ” which represents the point where the shallow edge of the capitellum starts. To infer “ β ” we make a basic geometric calculation based on the above assumptions, and we deduce that “ β ” is equal to 12 degrees. Now we calculate the residual distance the radial head must make to reach the critical point. 12 degrees is 0.2 radians leaving 1.8 radians for the radius to make from the original 2 radians. If we take the translation potential of the radial head following injury to the posterior capsule, this leaves us with 0.8 radians up to the critical point “ γ ” angle. 0.8 radian is the safe zone. This distance equals roughly a quarter of the functioning capitellum surface (Figure 2).




Figure 1: a. Sagittal MRI cut for the capittelum and the radial head in extension. The “ α ” angle represent 2 radians and the orange arrow cross at the critical point. b. Sagittal CT cut that demonstrate the safe zone wich equals roughly a quarter of the functional capitellar surface. “ γ ” angle is the angle between the blue arrow and orange arrow.




Figure 2: A simple 2D model for Radiocapitellar joint motion. a. 90 degrees elbow flexion. b. The elbow in neutral extension. c. one radian translation of the radial head following injury. d. Following “Osborne-Cotterill Lesion” of more than 0.8 radian, the radial head engages the lesion- “Mooji effect”.

3.1. This Safe Zone Could Be Breached in Several Scenarios

1. Fracture of the anterior articular surface of the radial head leaving less safe zone.
2. Classic “Osborne-Cotterill Lesion” with encroachment of the depression in the capitellum more anteriorly up to or passing the critical point. This could potentially lead to a large “ β ” angle. If “ β ” is equal to 57 degrees, then there is no safe zone, then, full extension, and supination with 1 radian translation will cause an engagement of the radial head on the capitellar depression- “Mooji effect”.
3. A combination between radial head fracture and “Osborne-Cotterill Lesion”. If we lose a third of the anterior radial head (0.66 radian) then, this will only leave little room for engagement (0.14 radian) [19,20].
4. Posterior comminution fracture of the capitellum or shear fracture of the capitellum.

4. Case Discussion

1.1. Case One

We present an 18-year-old patient who fell from a bicycle and

injured his right elbow. He was diagnosed with a fracture dislocation of the elbow (Figure 3a). The elbow was reduced in the ER, and a cast was applied (Figure 3b). A subsequent CT with 3D reconstruction revealed olecranon fracture with avulsion fracture of the Lateral ulnar collateral ligament (LUCL) and a posterior depression of the capitellum – “Osborne-Cotterill Lesion” (Fig 4.). The patient was offered surgery to fixate the fracture and stabilize the elbow joint. A posterior approach was utilized first, and the olecranon fracture was reduced and stabilized with locked plate (Figure 5a). Following this, a second lateral elbow approach with Kocher extension was utilized. LUCL was avulsed from its origin and there was a depression fracture on the posterior edge of the capitellum. At this stage, we feared that mere ligament stabilization will not be enough to prevent radial head engagement, thus, we proceeded to fill the depression with bone graft and a plate was contoured and utilized to stabilize the construct; Two anchors were used to reattach the LUCL (Figure 5b). Following the surgery, the elbow was splinted for a week, and then substituted with a dynamic elbow splint to allow early range of motion (Figure 6). 8 weeks following the surgery, the patient had full range of motion and was able to return fully to prior level of activities (Figure 7).




Figure 3: X-rays of the patient. a. fracture dislocation of right elbow. B. reduced elbow with posterior splint



Figure 4: 3D reconstruction of the injury following reduction




Figure 5.a: Fluoroscopic image following olecranon reduction and fixation with contoured plate. The elbow is unstable at this juncture. B. The “Osborne-Cotterill Lesion” was filled with bone graft and stabilized with contoured plate. Two anchors were used to reattach the LUCL. Following the procedure the elbow was stable in full range of motion.




Figure 6: X-ray images one week following the surgery. a. AP X-ray showing the posterior plate at the distal lateral part of the humerus. b. Lateral X-ray demonstrating the distal edge of the plate increasing the safe zone.




Figure 7: The patient after few weeks with full range of motion including full stable extension with supination.

1.2. Case Two

We present a 45-year-old patient who fell from a ladder and injured his left elbow. In the ER, he was diagnosed with a fracture of the capitellum and trochlea. A CT was done with 3D reconstruction to elucidate further the fracture and plan intervention (Figure 8). The 3D reconstruction revealed posterior capitellar comminution (Figure 9). A temporary splint was utilized for pain control, and the patient was admitted for surgical intervention. The next day, the patient underwent surgery with a lateral elbow approach and distal Kocher extension. The proximal incision with partial brachi-

oradialis release facilitated a deeper window for the trochlea. The trochlea was addressed first followed by the capitellum with reduction and provisional fixation with Kirchner wires and then with cannulated screws in the sagittal and coronal plains to maximize purchase and stability. A posterior fracture of the capitellar wall with comminution distally led us to add a posterior humeral plate advanced distally as close to the critical point to add stability and counteract the potential “Mooji effect” (Figure 10). The patient was allowed an early supervised range of motion 3 days following the surgery. The patient was discharged after 5 days to ambulatory rehabilitation and was lost to follow-up.




Figure 8: CT of the left elbow of the 2nd patient. a. The cut demonstrates the posterior lesion in the capitellum laterally. b. more medial cut shows the trochlear fracture.




Figure 9: 3D reconstruction of the fracture involving capitellum and trochlea with posterior comminution of trochlea inducing “Osborne-Cotterill Like -Lesion”.




Figure 10: Fluoroscopic and X-ray images of the patient left elbow following reduction and fixation with cannulated screws for the coronal fracture of the trochlea and capitellum and posterior contoured plate for the posterior comminution neutralizing the “Mooji effect”.

5. Discussion

We present a new term called the “Mooji effect, trying to explore the so-called “Osborne-Cotterill Lesion” further. We try to understand its impact and describe the best approach to address it to minimize complications and optimize outcome. We define the critical point of the capitellum as the most anterior point the radial head can reach before it dislocates and potentially engages a shallow area posteriorly. We termed the process of engagement the “Mooji effect” in reference to the author’s byname. The radial head can translate posteriorly up to the critical point. we figure the distance to be 1.8 times the radius of the capitellum. We assign the radial head a freedom of one radius translation in extreme position of extension and supination in the context of injury to the posterior capsule, leaving a mere 0.8 radius which is the safe zone. Based on our assumptions, the critical area can move further anteriorly encroaching on the safe zone up to a point that the radial head is inherently unstable in functional positions. This could be attributed to depression fracture as the radial head dislocates and imprints upon the capitellum or from comminuted fracture. We suggest that when the capitellum loses 0.25 of its posterior cartilage, then an osseous reconstruction is mandatory. When there is an additional fracture in the anterior radial head or the patient has joint laxity, then more liberal approach is warranted to address “Osborne-Cotterill Lesion”. While several authors introduced variant approaches to enhance osseous stability in the context of posterolateral instability, including reduction [10], bone grafting [13], osteochondral allograft transplantation [21], and even Prosthetic resurfacing [22], here we suggest a simple practical approach that helps reach stability with filling the defect with bone graft and utilizing a posterior plate to the capitellum to countereffect the “Mooji effect”.

United Prime Publications LLC., <https://acmcasereport.org/>

Our concept aligns with other biomechanical study that referred to radial head translation following posterolateral capsule injury that stated about 11 mm translation following injury to LUCL+ posterior capsule [12]. This in fact is half of the mean radial head diameter (one radian) in accordance with our assumption [23]. The plate not only protects the reduction or the lock bone graft, but its bulk and position prevent mechanically the “Mooji effect” and the radial head must translate further to skip the plate which increase the safe zone. This step should be contemplated whenever “Osborne-Cotterill Lesion” is found in the context of lateral elbow dislocation, fracture dislocation or complex elbow fractures. Other elements of the injury should be addressed including fractures and ligament avulsions. This will minimize the freedom of posterior translation of the radial head. Our theory resembles much the “On-Track/Off-Track” paradigm introduced by Itoi and associates [24,25], which discusses stability concept of the shoulder in the context of osseous defect in the glenoid and hill-Sachs lesion in the humeral head. Abduction and external rotation in the shoulder parallels extension and supination in the elbow and engagement happens easily with bipolar injury to the glenoid and the humerus as is the matter when the radial head loses anterior lip and the critical point of the capitellum moves anteriorly leading to “Mooji effect”. While off track lesions are treated with remplissage to the humeral head or Latarjet procedure to the glenoid [26], the “Osborne-Cotterill Lesion” leading to the “Mooji effect” is treated with bone graft and plating and the radial head fracture is either reduced and fixated or replaced. Interestingly, loss of more than 25% of the glenoid is treated with bony procedure like Latarjet procedure [26]. Other scenarios where we could encounter the “Mooji effect” and needs to be addressed is capitellar fractures with posterior comminution.

This could happen in the context of shear fractures of the capitellum elucidated through the Dubberley's classification of capitellar and/or trochlear fractures, specifically type B [27]. The posterior comminution, even with articular reduction could potentially lead to posterior engagement eventually, and it is addressed with bone grafting [28]; Furthermore, outcomes are inferior to type A without posterior comminution [29]. This is why it is better treated with additional posterior locked plate [30]. Another interesting option is shear capitellar fracture that leaves the radius engaged in without dislocation (Figure 11).



Figure 11: X-ray that demonstrates Capitellar fracture. (Bryan and Morrey Type I). The radial head is engaged in the fracture site.

6. Conclusion

We introduce an intuitive system to analyze the safe zone and help build a practical plan when “Osborne-Cotterill Lesion” is recognized. We encourage using CT scan in the sagittal plain and building 3D model to find the the anterior border of the lesion and see whether it encroaches upon the critical point and minimize the safe zone and beyond. If the safe zone is compromised and there is an accompanying fracture in the radial head further minimizing the room for maneuver, then moving back the critical zone is recommended by means of grafting and plating and not only ligament stabilization or reconstruction. We assign 25% safe zone and if it is violated then a bony procedure including plating, is recommended. Our study is not without shortcomings. It only demonstrates two cases with one case lost to follow-up. Our recommendations are based on multiple hypotheses that needs to be further explored in laboratory conditions and clinical set-up. More evidence is needed to collaborate our assumptions. But taking into consideration the rarity of this injury and the scarce literature that is based on case reports and small series, then we strongly advise re-evaluating the “Osborne-Cotterill Lesion” and adopting the concept of “Mooji effect” when treating similar injuries.

References

1. Robinson PM, E Griffiths. Simple elbow dislocation. *Shoulder & Elbow*. 2017; 9(3): 195-204.
2. de Haan J. Simple elbow dislocations: a systematic review of the literature. *Archives of orthopaedic and trauma surgery*. 2010; 130: 241-249.
3. Hackl M. The Treatment of Simple Elbow Dislocation in Adults: A Systematic Review and Meta-analysis. *Deutsches Ärzteblatt International*. 2015; 112(18): 311.
4. Geyer S. Patient satisfaction, joint stability and return to sports following simple elbow dislocations: surgical versus non-surgical treatment. *Archives of orthopaedic and trauma surgery*. 2023; 143(3): 1481-1489.
5. Osborne G, P Cotterill. Recurrent dislocation of the elbow. *J Bone Joint Surg Br*. 1966; 48(2): 340-6.
6. Somford MP. The Osborne-Cotterill lesion: how an eponymous term arose and evolved. *JBJS*. 2019; 101(16): e81.
7. Jeon IH. Osborne-cotterill lesion: an osseous defect of the capitellum associated with instability of the elbow. *American Journal of Roentgenology*. 2008; 191(3): 727-729.
8. Lee JH, TK Lim. We all know about the Osborne-Cotterill lesions, but we still don't know how to treat them. *Clinics in Shoulder and Elbow*. 2023; 26(3): 220.
9. Yano K, Y Kaneshiro. Acute elbow trauma similar to Osborne-Cotterill lesion, capitellar impaction fracture associated with posterolateral rotatory instability: A case report and literature review. *Journal of Orthopaedic Science*. 2021; 26(6): 1152-1155.
10. Vargas DG. Osborne-Cotterill lesion a forgotten injury: review article and case report. *Clinics in shoulder and elbow*. 2020; 23(1): 27.
11. Jeon IH. Surgical treatment and clinical implication for posterolateral rotatory instability of the elbow: Osborne-Cotterill lesion of the elbow. *Journal of Trauma and Acute Care Surgery*. 2011; 71(3): E45-E49.
12. Edwards DS. The contribution of the posterolateral capsule to elbow joint stability: a cadaveric biomechanical investigation. *Journal of shoulder and elbow surgery*. 2018; 27(7): 1178-1184.
13. Shekhibi A, W Reichert. Engaging Osborne-Cotterill lesion as a cause of irreducible posterior dislocation of the radiocapitellar and ulnohumeral joints: A case report. *Journal of Orthopaedic Reports*. 2022; 1(4): 100066.
14. Alcid JG, CS Ahmad, TQ Lee. Elbow anatomy and structural biomechanics. *Clinics in sports medicine*. 2004; 23(4): 503-517.
15. Griswold BG. Radius of curvature of the radial head matches the capitellum: a magnetic resonance imaging analysis. *JSES international*. 2023; 7(4): 668-672.
16. Dunning CE. Quantifying translations in the radiohumeral joint: application of a floating axis analysis. *Journal of biomechanics*. 2003; 36(8): 1219-1223.
17. Kang CN, CH Choi. The Normal Humero-Capitellar Angle in Children; Baumann Angle and Lateral Humero-Capitellar Angle. *Emj*. 1994; 17(4): 383-390.

18. Rotman D. The posterolateral ligament of the elbow: anatomy and clinical relevance. *Journal of Shoulder and Elbow Surgery*. 2024; 33(3): 573-582.
19. Nalbantoglu U. Capitellar cartilage injuries concomitant with radial head fractures. *The Journal of hand surgery*. 2008; 33(9): 1602-1607.
20. Claessen FM. Outcomes of concomitant fractures of the radial head and capitellum: the “kissing lesion”. *Journal of Hand and Microsurgery*. 2016; 8(02): 100-105.
21. Lee SJ. Elbow osteochondral allograft transplantation and lateral ulnar collateral ligament repair with internal brace: a case report. *JBJS Case Connector*. 2022; 12(1): e21.
22. Rotman D, JR Lievano. Prosthetic resurfacing of engaging posterior capitellar defects in recurrent posterolateral rotatory instability of the elbow. *Clinics in Shoulder and Elbow*. 2023; 26(3): 287-295.
23. Guitton TG, HJ Van Der. Quantitative measurements of the volume and surface area of the radial head. *The Journal of hand surgery*. 2010; 35(3): 457-463.
24. Yamamoto N. Contact between the glenoid and the humeral head in abduction, external rotation, and horizontal extension: a new concept of glenoid track. *Journal of shoulder and elbow surgery*. 2007; 16(5): 649-656.
25. Di Giacomo. Burkhart, Evolving concept of bipolar bone loss and the Hill-Sachs lesion: from “engaging/non-engaging” lesion to “on-track/off-track” lesion. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2014; 30(1): 90-98.
26. Itoi E. ‘On-track’and ‘off-track’shoulder lesions. *EFORT Open Reviews*. 2017; 2(8): 343-351.
27. Teng L, G Zhong. Surgical treatment of comminuted coronal shear fracture of distal humerus. *Orthopaedic Surgery*. 2020; 12(5): 1439-1447.
28. Tarallo L. Surgical tips and tricks for coronal shear fractures of the elbow. *Archives of Orthopaedic and Trauma Surgery*. 2021; 141: 261-270.
29. Marinelli A. Does the presence of posterior comminution modify the treatment and prognosis in capitellar and trochlear fractures? Study performed on 45 consecutive patients. *Injury*. 2018; 49: S84-S93.
30. Gao X. Clinical outcomes of capitellar fractures with posterior comminution treated with Herbert screws combined with metacarpal locking plates. *BMC Musculoskeletal Disorders*. 2023; 24(1): 937