

# Pulmonary Rehabilitation: A Cornerstone in Post-Lung Cancer Surgery Care

Zeng Yanyan<sup>1</sup>, Yang Yanli<sup>1</sup>, Zhang Guangying<sup>1</sup>, Li Mingyang<sup>1</sup>, Xin Yangyang<sup>1</sup> and Xu Hui<sup>2\*</sup>

<sup>1</sup>Department of Thoracic Surgery Medical Center, Weifang people's hospital; the first affiliated hospital of shandong second medical university. 261041, Weifang, Shandong, P.R. China

<sup>2</sup>Department of Nursing, Weifang people's hospital; the first affiliated hospital of shandong second medical university. 261041, Weifang, Shandong, P.R. China

## \*Corresponding Author:

Xu Hui, MD, Department of Nursing, Weifang people's hospital; the first affiliated hospital of shandong second medical university, 261041, Weifang, Shandong, P.R. China

**Received:** 13 Apr 2025

**Accepted:** 25 Apr 2025

**Published:** 28 Apr 2025

**J Short Name:** ACMCR

**Copyright:** ©2025 X Hui. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

**Keywords:** Pulmonary Rehabilitation; Lung Cancer; Postoperative Complications; Exercise; Quality of Life

**Citation:** X Hui. Pulmonary Rehabilitation: A Cornerstone in Post-Lung Cancer Surgery Care. *Anna Clin Med Case Rep*. 2025; 14(11): 1-4

## 1. Abstract

Lung cancer remains a leading cause of cancer-related mortality globally, and surgical resection is a primary treatment for early-stage disease. Postoperative pulmonary complications (PPCs) significantly impact patient outcomes and recovery. Pulmonary rehabilitation (PR) has emerged as a critical intervention to mitigate the adverse effects of lung resection and enhance overall well-being. This review synthesizes current evidence on PR for post-lung cancer surgery patients, encompassing its benefits, implementation challenges, and future research directions. While existing studies demonstrate the efficacy of PR in improving pulmonary function, exercise capacity, and quality of life, the optimal program design and long-term outcomes require further investigation. Large-scale randomized controlled trials, standardized PR protocols, and patient-centered research are essential to advance the field and optimize PR delivery for post-lung cancer patients.

## 2. Introduction

Lung cancer, a malignant neoplasm originating from the bronchial mucosa or glands, poses a significant global health burden [1]. As reported by the World Health Organization (WHO), it remains the leading cause of cancer-related mortality in 87 countries, with a staggering 1.79 million deaths attributed to the disease in 2020 alone [2]. This underscores the urgent need for comprehensive strategies to prevent, detect, and manage lung cancer. Despite advancements in diagnostic and therapeutic modalities, surgical resection remains the cornerstone of curative treatment for early-stage non-small cell lung cancer (NSCLC) [3]. However, pulmonary resection is associated with substantial morbidity, including a heightened risk of postoperative pulmonary complications (PPCs) such as pneumonia, respiratory failure, and atelectasis [4, 5]. These complications significantly impact patient outcomes, prolong hospital stays, increase healthcare costs, and diminish quality of life [6]. To mitigate the adverse sequelae of lung resection and optimize patient recovery, pulmonary rehabilitation (PR) has emerged as a critical component of postoperative care. A growing body of evidence supports the efficacy of PR in improving pulmonary function, exercise tolerance, and overall well-being in patients with lung cancer [7, 8]. By addressing physical, psychological, and educational needs, PR aims to reduce the incidence and severity of PPCs, facilitate a faster return to independence, and enhance long-term survival [9]. This review comprehensively examines the current state of knowledge regarding PR for post-lung cancer surgery patients, highlighting its benefits, challenges, and future directions.

## 3. Problems After Lung Cancer Surgery

Surgical resection of lung tissue, while curative in early-stage NSCLC, inevitably results in physiological alterations that

compromise pulmonary function. This functional impairment manifests as dyspnea, reduced exercise capacity, and diminished quality of life [10]. Moreover, the invasive nature of lung surgery predisposes patients to a range of postoperative complications, including pneumonia, respiratory failure, and pulmonary embolism, which can significantly prolong hospital stays, increase healthcare costs, and elevate mortality rates [11, 12]. A retrospective analysis revealed that the incidence of postoperative complications within 30 days of lung resection can be as high as 58%, with a mortality rate of 3.6% during the same period [13]. These findings underscore the substantial burden of postoperative morbidity and mortality associated with lung cancer surgery and highlight the imperative for effective interventions, such as pulmonary rehabilitation, to optimize patient outcomes.

## 4. Concepts and Training Methods Related to Pulmonary Rehabilitation

### 4.1. Concepts Related to Pulmonary Rehabilitation

Pulmonary rehabilitation (PR) is a multidisciplinary, evidence-based intervention designed to improve the physical and psychological well-being of individuals with chronic respiratory conditions [11,12]. The concept of PR emerged in the mid-20th century, with the American College of Chest Physicians introducing the term in 1974 [11]. Since then, the field has evolved significantly, with a growing body of evidence supporting its efficacy. A seminal consensus statement by the American Thoracic Society (ATS) and the European Respiratory Society (ERS) in 2013 defined PR as a comprehensive program that includes but is not limited to exercise training, education, and behavior modification [12]. This program is tailored to the individual patient based on a thorough assessment of their respiratory condition, functional status, and psychosocial needs. The overarching goal of PR is to optimize functional status, reduce symptoms, and enhance the overall quality of life for individuals with chronic respiratory diseases. Beyond its physical benefits, PR plays a crucial role in addressing the psychological and social impact of chronic respiratory conditions. By providing education and support, PR empowers patients to manage their disease, adhere to treatment plans, and participate actively in their healthcare. Importantly, PR is applicable across the spectrum of respiratory diseases, including various stages of lung cancer, demonstrating its versatility and potential to improve outcomes for a wide range of patients. The National Institute for Health and Clinical Excellence (NICE) guidelines have further underscored the importance of PR in reducing mortality and pulmonary complications, while improving lung function in patients with chronic respiratory diseases [13,14]. These guidelines highlight the cost-effectiveness of PR and emphasize its role as a cornerstone of comprehensive care for individuals with respiratory conditions.

## 4.2. Pulmonary Rehabilitation Training Methods and Evaluation Indicators

### 4.2.1. Training Methods

At present, the methods of postoperative pulmonary rehabilitation training adopted in clinical practice vary. Because pulmonary rehabilitation training contains a lot of content, many researchers develop personalized training methods based on the specific conditions of patients and research purposes. Pulmonary rehabilitation training mainly includes exercise training, health education, nutritional support and psychological education. Its core is exercise training. According to the different parts of the exercise, exercise training can be divided into upper limb training and lower limb training. Upper limb training methods include finger wall climbing exercises, elbow flexion and extension exercises, upper limb weight lifting, etc.; lower limb training methods include walking, climbing stairs, jogging and cycling, etc. According to the purpose of training, exercise training can be divided into: endurance training, interval training, resistance training, respiratory muscle exercise and balance training [15]. Endurance training is recommended to be performed at least 3 times a week for 8 weeks. The most common endurance training mode is walking (ground walking and stationary cycling); interval training consists of high-intensity exercise (20 s to 30 min) and rest or lower-intensity training (30 s to 30 min), including cycling and walking [16]. Interval training during cycling or walking avoids the need for long rest periods, thereby minimizing interruptions to the exercise process and can be completed with less dyspnea and leg fatigue [16-18]. Resistance training refers to the activation of local muscle groups in relatively high-load repetitive weightlifting exercises [15]. Resistance training is recommended to be performed 1 to 3 sets of 8 to 12 repetitions per set 2 or 3 days a week [19], including resistance training of the upper and lower limbs. Common respiratory muscle exercises include pursed lip/abdominal breathing, yoga gymnastics, balloon blowing, etc. [20]. There is no consensus on the optimal duration of respiratory muscle training, but most studies provide a total exercise duration of 30 to 60 minutes per day, usually 2 to 3 times, and an average of 3 to 7 days per week [21]. Balance training is becoming an adjunct to traditional PR training programs [15]. Balance training includes circuit training such as standing exercises and gait transitions.

### 4.2.2. Evaluation Indicators

**1) Functional exercise capacity:** The 6-minute walk test (6 MWT) is used to assess functional and motor capacity. The test is performed according to published guidelines and the better of 16 and 2 attempts is recorded [22]. A 30-meter corridor is used and the patient is instructed to walk as far as possible in 6 minutes. Standardized encouragement is given at the end of each minute. Oxygen saturation and pulse rate are continuously monitored throughout the test, and the modified Borg volume is used immediately after the test to quantify the degree of dyspnea and leg fatigue.

**2) Skeletal Muscle Strength:** Handgrip strength (HF) and quadriceps strength (QF) were measured as indicators of skeletal muscle strength. 1 HF was assessed using a dynamometer. The patient was required to squeeze the dynamometer with maximal force for at least 5 seconds. After 1 exercise attempt, HF was tested twice with at least 60 seconds of rest between each attempt. The highest value of the 2 attempts for both hands was recorded (kg). Cardiac strength was measured for both limbs to consider the potential impact of postoperative pain on the operated side. During the test, the patient stood upright with his feet hip-width apart, elbows fully extended, holding the dynamometer, wrist in neutral position, and index finger flexed 90°. 2 Quadriceps strength was assessed as the peak force generated by the leading leg during a seated maximal isometric knee extension exercise using a handheld dynamometer with a strap, and the highest value of at least 2 maneuvers was recorded and expressed in kilograms feet (kgf).

**3) Respiratory Function:** The respiratory function of patients was tested by spirometry, including forced vital capacity (FVC), forced expiratory volume in 1 second (FEV1), percentage of forced expiratory volume in 1 second to predicted value (FEV1%), and ratio of forced expiratory volume in 1 second to forced vital capacity (FEV1/FVC) [23].

**4) Quality of Life:** The core quality of life scale (EORTC QLQ-C30) was used to assess the quality of life of patients. The scale has 5 dimensions (physical, emotional, role, cognitive, and social) and 30 items. Each item is scored on a scale. The higher the overall score, the higher the quality of life of the patient.

## 5. Current Status of Pulmonary Rehabilitation (PR) After Lung Cancer Surgery

Pulmonary rehabilitation (PR) is a minimally invasive treatment intervention that provides patients with a variety of tools to improve respiratory symptoms and quality of life [24,25]. Studies have shown that exercise training can improve the exercise capacity and quality of life of lung cancer patients undergoing surgery [26]. Xu Haijiang et al. [27] believed that the combination of osimertinib and pulmonary rehabilitation training can improve the lung function of patients with non-small cell lung cancer (NSCLC) who underwent radical resection of lung cancer, reduce their postoperative morbidity, and thus improve their quality of life, which is conducive to reducing the medical burden on patients and society. A meta-analysis [28] included 8 randomized controlled trials to explore the effects of aerobic exercise and resistance training on patients after lung cancer surgery. The study showed that exercise training can improve the exercise capacity and quadriceps strength of patients after lung resection of non-small cell lung cancer. In a randomized controlled trial by Scott et al. [29], lung cancer survivors with poor cardiopulmonary fitness (CFR) were included in 48 three-week continuous supervised courses, which included aerobic training (AT), combined training (CT) and resistance training (RT). The study showed that AT and CT significantly improved the peak maximum oxygen uptake (Vo2) of lung cancer survivors. At the same time, multiple studies have shown that pulmonary rehabilitation can also improve quality of life, forced expiratory volume in 1 second and forced vital capacity, and reduce lung cancer complications and mortality [30-33]. In addition, most lung cancer patients have nutritional problems to varying degrees. PR provides an opportunity to educate patients to maintain a proper diet [24]. The combination of increased calorie intake, exercise, and in some cases the use of anabolic steroids may lead to significant weight gain. Due to calorie intake, more high-protein foods need to be eaten. Protein is broken down into amino acids by the human body. These amino acids are absorbed by the body and can be synthesized into muscle through proper exercise, thereby increasing weight. In addition, since lung cancer patients have a large nutritional demand after surgery, standardized nutritional intervention can reduce the degree of protein breakdown in the patient's body, avoid deterioration of nutritional status, and achieve the purpose of maintaining and gradually improving nutritional status [34].

## 6. Limitations of Postoperative Pulmonary Rehabilitation

While the extant literature demonstrates the potential benefits of pulmonary rehabilitation (PR) in enhancing lung function and exercise capacity among post-lung cancer surgery patients, several limitations persist.

## 7. Methodological Challenges

### 7.1. Heterogeneity of Study Designs

The existing body of evidence is characterized by heterogeneity in study designs, including variations in participant selection criteria, intervention modalities, and outcome measures. This heterogeneity impedes the establishment of robust conclusions regarding the optimal PR program and its effectiveness.

**Small Sample Sizes:** Many studies suffer from limited sample

sizes, which can reduce statistical power and limit the generalizability of findings. Larger-scale randomized controlled trials (RCTs) are needed to provide more definitive evidence.

## 7.2. Lack of Long-Term Outcomes

Most studies focus on short-term outcomes, such as changes in pulmonary function and exercise capacity. There is a dearth of research investigating the long-term effects of PR on quality of life, survival, and healthcare utilization.

## 7.3. Adherence Challenges

Ensuring patient adherence to PR programs remains a significant obstacle. Factors such as physical limitations, comorbidities, psychological distress, and lack of social support can hinder patient engagement.

## 7.4. Clinical Implementation Gaps

**Standardized Protocols:** The absence of standardized PR protocols hampers the ability to compare outcomes across studies and implement evidence-based practices consistently.

## 7.5. Access and Delivery

Geographic disparities in access to PR services and variations in the delivery of care models contribute to inconsistent outcomes.

## 7.6. Integration with Cancer Care

Integrating PR into the continuum of cancer care is essential but remains a challenge. Effective collaboration between surgeons, oncologists, and rehabilitation specialists is crucial for optimal patient management.

## 7.7. Resource Constraints

Limited resources, including personnel, facilities, and funding, can impede the widespread implementation of PR programs.

## 8. Future Research Directions

Addressing these limitations requires a concerted effort to conduct well-designed RCTs with larger sample sizes, investigate long-term outcomes, and develop standardized PR protocols. Moreover, research focused on identifying patient-specific factors influencing PR adherence and developing strategies to enhance engagement is warranted. By overcoming these challenges, the field can advance the implementation of effective PR programs for post-lung cancer surgery patients.

## 9. Summary and Outlook

Pulmonary rehabilitation (PR) has emerged as a critical component of post-lung cancer surgery care. Accumulating evidence underscores its efficacy in improving pulmonary function, exercise tolerance, and quality of life, thereby enhancing patient outcomes and reducing the burden of postoperative complications. However, the current body of evidence on PR for post-lung cancer patients remains limited. While existing studies have demonstrated the benefits of PR, the optimal program design, including intensity, duration, and components, requires further investigation. Additionally, the long-term effects of PR on patient outcomes, such as survival and recurrence, warrant exploration. To advance the field of pulmonary rehabilitation in post-lung cancer care, several key areas warrant future research: Large-scale randomized controlled trials: Conducting rigorously designed randomized controlled trials with adequate sample sizes is essential to establish the efficacy and cost-effectiveness of PR programs.

### 9.1. Standardized PR Protocols

Developing standardized PR protocols tailored to the specific needs of post-lung cancer patients will facilitate the implementation and evaluation of PR programs across different healthcare settings.

### 9.2. Longitudinal Studies

Investigating the long-term impact of PR on patient outcomes,

including quality of life, functional status, and survival, is crucial to inform clinical practice and healthcare policy.

### 9.3. Patient-Reported Outcomes

Incorporating patient-reported outcome measures into future research will provide valuable insights into the patient experience and the impact of PR on their overall well-being.

### 9.4. Multidisciplinary Collaboration

Fostering collaboration among pulmonologists, surgeons, rehabilitation specialists, and other healthcare providers is essential for optimizing PR delivery and outcomes. By addressing these research priorities, the field can move towards evidence-based guidelines for PR in post-lung cancer care, ultimately improving patient outcomes and enhancing the quality of life for lung cancer survivors.

## References

1. Park YS, LeeJ, Keum B, Oh BM. Feasibility of an Eight-Week Outpatient-Based Pulmonary Rehabilitation Program for Advanced Lung Cancer Patients Undergoing Cytotoxic Chemotherapy in Korea. *Thoracic Cancer*. 2018; 9: 1069-1073.
2. LIU Zongchao, LI Zhexuan, ZHANG Yang. Interpretation of the 2020 Global Cancer Statistics Report[J]. *Electronic Journal of Comprehensive Cancer Therapy*. 2021; 7(2): 1-14.
3. PENG Bo, HUANG Yubui, HUANG Houqiang. Systematic evaluation and meta-analysis of the effects of postoperative pulmonary rehabilitation on lung cancer patients. *Modern Medicine and Health*. 2021; 37(22): 3836-3880.
4. Wang L, Yu M, Ma Y, Tian R. Effect of Pulmonary Rehabilitation on Postoperative Clinical Status in Patients with Lung Cancer and Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-Analysis. *Evidence-Based Complementary and Alternative Medicine*. 2022; 4133237.
5. ZHAO Ying, LU Jia. Study on the significance of pulmonary rehabilitation training for postoperative chemotherapy patients with lung cancer[J]. *China Practical Medicine*. 2023; 18(5): 171-173.
6. Xu Huan, Zhu Xingxing, Liu Zhihao. Effects of early postoperative pulmonary rehabilitation training on postoperative pulmonary function and quality of life of lung cancer patients[J]. *Zhejiang Medicine*. 2020; 42(22): 2443-2445.
7. Liu Y, Wu A, Li X. Retrospective Analysis of Eleven Gene Mutations, PD-L1 Expression and Clinicopathological Characteristics in Non-Small Cell Lung Cancer Patients. *Asian Journal of Surgery*. 2022; 45: 367-375.
8. Deng WJ, Zhou WD. The clinical value of rehabilitation training on lung function and survival quality of postoperative lung cancer patients. *J Chinese Practical Medicine*. 2023; 18(3): 159-163.
9. Deng J, Li R. The Effect of Rehabilitation Training on Postoperative Patients with Lung Cancer: A Systematic Review and Meta-Analysis. *Asian Journal of Surgery*. 2022; 45: 1968-1970.
10. Andalib A, Ramana-Kumar AV, Bartlett G, Franco EL. Influence of Postoperative Infectious Complications on Long-Term Survival of Lung Cancer Patients: A Population-Based Cohort Study. *Journal of Thoracic Oncology*. 2013; 8: 554-561.
11. Meng S. Research progress of pulmonary rehabilitation[J]. *Zhonghua Medical Information Herald*. 2007; 20-22.
12. Spruit MA, Singh SJ, Garvey C, ZuWallack R. An Official American Thoracic Society/European Respiratory Society Statement: Key Concepts and Advances in Pulmonary Rehabilitation. *American Journal of Respiratory and Critical Care Medicine*. 2013; 188: e13-e64.
13. Poston G, Tait D, Beattie R, Byrne C, Chapman J. National Institute for Health and Clinical Excellence: The Diagnosis and

- Management of Colorectal Cancer: Full Guideline. 2011.
14. Jamison DT, Gelband H, Horton S, Jha P. Disease Control Priorities: Improving Health and Reducing Poverty. 3rd Edition, The International Bank for Reconstruction and Development/The World Bank, Washington DC. 2017.
  15. Lee AL, Holland AE. Time to Adapt Exercise Training Regimens in Pulmonary Rehabilitation-A Review of the Literature. *International Journal of Chronic Obstructive Pulmonary Disease*. 2014; 9: 1275-1288.
  16. Kortianou EA, Nasis IG, Spetsiotti ST. Effectiveness of Interval Exercise Training in Patients with COPD. *Cardiopulmonary Physical Therapy Journal*. 2010; 21: 12-19.
  17. Vogiatzis I, Nasas S, Roussos C. Interval Training as an Alternative Modality to Continuous Exercise in Patients with COPD. *European Respiratory Journal*. 2002; 20: 12-19.
  18. Sabapathy S, Kingsley RA, Schneider DA. Continuous and Intermittent Exercise Responses in Individuals with Chronic Obstructive Pulmonary Disease. *Thorax*. 2004; 59: 1026-1031.
  19. American College of Sports Medicine American College of Sports Medicine Position Stand Progression Models in Resistance Training for Healthy Adults. *Medicine & Science in Sports & Exercise*. 2009; 41: 687-708.
  20. QIN Mengxia, QIAN Yan, CHEN Ying. Progress of pulmonary rehabilitation in the treatment of lung cancer patients. *Journal of Nursing*. 2019; 34(10): 101-104.
  21. American Thoracic Society/European Respiratory Society. ATS/ERS Statement on Respiratory Muscle Testing. *American Journal of Respiratory and Critical Care Medicine*. 2002; 166: 518-624.
  22. Oikawa M, Hanada M, Nagura H, Tsuchiya T. Factors Influencing Functional Exercise Capacity after Lung Resection for Non-Small Cell Lung Cancer. *Integrative Cancer Therapies*. 2020; 19: 1-8.
  23. Muhua, Li Yufeng. Effects of pulmonary rehabilitation exercise on postoperative respiratory function recovery and exercise tolerance in patients undergoing radical lung cancer surgery. *J Clinical Medicine Research and Practice*. 2022; 7(8): 168-170.
  24. Rivas-Perez, H. and Nana-Sinkam, P. Integrating Pulmonary Rehabilitation into the Multidisciplinary Management of Lung Cancer: A Review. *Respiratory Medicine*. 2015; 109: 437-442.
  25. LIU Xiaohong, ZHANG Qiong. Effects of pulmonary rehabilitation training nursing on lung function and quality of life of postoperative chemotherapy patients with lung cancer[J]. *Wisdom Health*. 2021; 7(6): 110-198.
  26. Reeve J, Stiller K, Nicol K, McPherson KM. A Postoperative Shoulder Exercise Program Improves Function and Decreases Pain Following Open Thoracotomy: A Randomised Trial. *Journal of Physiotherapy*. 2010; 56: 245-252.
  27. Xu H, Guo R, Yang Y. Effects of Osimertinib Combined with Pulmonary Rehabilitation and Health Care Training on Pulmonary Function, Complications, and Quality of Life in Patients after Radical Resection of Lung Cancer. *Frontiers in Public Health*. 2022; 10.
  28. Cavalheri V, Burtin C, Formico VR, Nonoyama ML. Exercise Training Undertaken by People within 12 Months of Lung Resection for Non-Small Cell Lung Cancer. *The Cochrane Database of Systematic Reviews*. 2019; 6: CD009955.
  29. Scott JM, Thomas SM, Herndon JE, Douglas PS. Effects and Tolerability of Exercise Therapy Modality on Cardiorespiratory Fitness in Lung Cancer: A Randomized Controlled Trial. *Journal of Cachexia, Sarcopenia and Muscle*. 2021; 12: 1456-1465.
  30. Tiep B, Sun V, Koczywas M, Kim J. Pulmonary Rehabilitation and Palliative Care for the Lung Cancer Patient. *Journal of Hospice & Palliative Nursing*. 2015; 17: 462-468.
  31. Vagvolgyi A, Rozgonyi Z, Kerti M, Agathou G, Vadasz P. Effectiveness of Pulmonary Rehabilitation and Correlations in between Functional Parameters, Extent of Thoracic Surgery and Severity of Post-Operative Complications: Randomized Clinical Trial. *Journal of Thoracic Disease*. 2018; 10: 3519-3531.
  32. Choi J, Yang Z, Lee J, Lee JH, Kim HK. Usefulness of Pulmonary Rehabilitation in Non-Small Cell Lung Cancer Patients Based on Pulmonary Function Tests and Muscle Analysis Using Computed Tomography Images. *Cancer Research and Treatment*. 2022; 54: 793-802.
  33. Mao X, Ni Y, Niu Y, Jiang L. The Clinical Value of Pulmonary Rehabilitation in Reducing Postoperative Complications and Mortality of Lung Cancer Resection: A Systematic Review and Meta-Analysis. *Frontiers in Surgery*. 2021.
  34. Mele MC, Rinninella E, Cintoni M, Pulcini G. Nutritional Support in Lung Cancer Patients: The State of the Art. *Clinical Lung Cancer*. 2021; 22: e584-e594.