

Phasix™ non-ST Mesh for Giant Paraesophageal Hernias - The Way of the Future?

Aldenb Lorenzo^{1*}, Sarah Mahmood^{1,2}, Kevin Jia¹, Xin Lin², David Goltsman¹ and Christophe R Berney^{1,2}

¹Department of Surgery, Bankstown-Lidcombe Hospital, Sydney, Australia

²Faculty of Medicine, University of New South Wales, Sydney, Australia

*Corresponding Author:

Aldenb Lorenzo, BSE, MD, FRACS, Department of Surgery, Bankstown-Lidcombe Hospital, Sydney, Australia

Received: 13 Sep 2025

Accepted: 20 Sep 2025

Published: 22 Sep 2025

J Short Name: ACMCR

Copyright:

©2025 A Lorenzo. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Keywords: Hiatus Hernia; Phasix Mesh; Laparoscopic Hiatus Hernia Repair

Citation: A Lorenzo. Phasix™ non-ST Mesh for Giant Paraesophageal Hernias - The Way of the Future?. *Anna Clin Med Case Rep*. 2025; 15(11): 1-5

1. Abstract

1.1 Background

Mesh reinforcement can be used as an adjunct in giant paraesophageal hernias repairs. The use of synthetic mesh is associated with significant morbidity whereas biologic mesh is associated with higher rates of recurrence. We aim to assess the safety and effectiveness of biosynthetic absorbable Phasix™ non-ST mesh in the repair of giant paraesophageal hernias.

1.2. Methods

A cohort study of all patients presenting to Bankstown-Lidcombe hospital for elective laparoscopic mesh repair of giant paraesophageal hernias was prospectively recorded. Preoperative investigations included gastroscopy and CT-scan of the chest and abdomen. Phasix™ non-ST mesh was used in all cases. Clinical follow ups were scheduled at 2- and 6-weeks, 4- and 12-months post-operative. All patients underwent yearly postoperative CT-scan and gastroscopy. Primary endpoint was endoscopic and/or radiological recurrence, and secondary endpoints included length of hospital stay, morbidity, mortality, and symptom recurrence.

1.3. Results

Thirty-two patients were included. Dyspnoea (62.5%) and dysphagia (53.1%) were the most common symptoms. Twenty (62.5%) had type III and five (15.6%) had type IV paraesophageal hernias. Median length of stay was 3 days (range 2-7) and only minor postoperative complications were recorded in two patients (6.3%). The median follow-up time was 26 months (range 12-53). No mesh-related complications were recorded and 30 patients (93.8%) were symptom-free. There was only one endoscopic and radiological recurrence found in one patient at 18 months post-surgery.

1.4. Conclusion

Phasix™ non-ST mesh reinforcement of the oesophageal hiatus is feasible with satisfactory symptoms improvement and no adverse outcomes. Further RCTs are required to investigate long-term efficacy.

2. Introduction

Hiatus hernias are relatively common findings in the adult population and often found incidentally on gastroscopy or imaging. We classify hiatus hernias into sliding (type I) and paraesophageal (types II, III and IV) hernias, with type I hernia being characterised by the displacement of the gastro-oesophageal junction (GOJ) above the diaphragm whilst the fundus remains below. Paraesophageal hernias (type II-IV) are considered true hernias as they possess a hernia sac and are characterised by the gastric fundus being displaced upwards through a defect in the phreno-oesophageal membrane, with

increasing abdominal contents protruding up into the defect. These paraesophageal hernias comprise 5-10% of all hiatal defects [1]. Patients can be asymptomatic, but often exhibit a broad spectrum of clinical symptoms which can include dyspepsia and reflux, epigastric or chest pain, dysphagia, nausea, regurgitation, shortness of breath, chronic cough, sore throat, and iron deficiency anaemia [2]. Patient may also present with severe complications secondary to paraesophageal hernias which can include gastrointestinal bleeding, pulmonary aspiration, gastric outlet obstruction, or gastric volvulus. Elective surgical repair is generally indicated in patients with symptomatic paraesophageal hernias but may be also influenced by other factors including patient age and the severity of existing co-morbidities. Although evidence suggests that there is no indication for intervention in those who are asymptomatic, there is a >10% per year risk to becoming symptomatic [3]. Furthermore, the annual probability of emergency surgery for paraesophageal hernias in the adult population is around 1%, with a lifetime risk of approximately 18% at 65 years of age [4]. Whilst the mainstay of modern paraesophageal hernia repair is laparoscopic, there remains considerable disparities in the techniques employed including variation in method of hiatal closure and choice of fundoplication, usage and mesh selection. Recurrence rates quoted in literature vary widely, ranging from 4%-66% [5-7]. The severity of recurrence can vary, with the majority being asymptomatic recurrences detected on imaging, whilst others clinically symptomatic requiring re-intervention. Given the potentially high recurrence rates, mesh reinforcement can be considered to circumvent this, however it comes with its own inherent risks. Indeed, permanent synthetic meshes have demonstrated long-term risks of mesh erosion and oesophageal stricture, whereas biological meshes have a hypothetical higher risk for recurrence as they don't integrate well due to limited foreign body response. However, two recent systematic reviews of 7 RCTs demonstrated recurrence rates may be equivocal in both mesh groups (whether synthetic or biological) and could not show any real benefit over simple suture repair. [8,9] Conversely, when looking at pooled data from six RCTs and 13 observational studies, Rajkomar et al. [8,9]. Established a significant reduction in large hiatal hernia recurrence rates with use of mesh [10]. In recent years, a slowly resorbable biosynthetic mesh has been introduced in Australia. Phasix™ non-ST mesh, or poly-4-hydroxybutyrate (P4HB), is a knitted monofilament mesh that fully resorbs within 12-18 months via hydrolysis. Phasix™ meshes (ST or non-ST) have been widely used worldwide in ventral and incisional hernia repairs. Other biosynthetic or absorbable meshes available include the Gore Bio-A® which has a significantly quicker resorption time of 6-8 months. Those absorbable meshes may reduce the risk of erosion, but hypothetically with the potential long-term cost of hernia recurrence. At present, mesh usage in the

hiatus remains less than 20% in Australia [11]. Due to limited data availability, questions remain as to whether biosynthetic meshes should be routinely used in laparoscopic large hiatus hernia repairs, to improve long-term efficacy (reduced recurrence and complication rates), as well as quality-of-life. Our observational single-institution prospective cohort study aims to assess the feasibility and safety of Phasix™ non-ST mesh with atraumatic fibrin glue fixation (Tisseel) during laparoscopic repair of large sliding hiatus and paraoesophageal hernias.

3. Methods

3.1. Study Design

We conducted an observational single-centre prospective cohort study. All adult patients (aged ≥ 18 years) admitted to Bankstown-Lidcombe Hospital under the care of one of the co-authors (CB) for elective laparoscopic repair of large (>7 cm) sliding (type I) hiatus or paraoesophageal (types II to IV) hernias, using biosynthetic Phasix™ non-ST mesh reinforcement, from September 2019 to March 2022, were included. Data was entered into a prospectively maintained database. We recorded patient demographics, pre-operative symptomatology and ASA score. All patients underwent routine pre-operative clinical assessment, including CT chest and gastroscopy. Selected patients also underwent pre-operative transthoracic echocardiogram (TTE) motivated by chest pain and/or dyspnoea. Our primary outcome of interest was endoscopic recurrence, defined as a recurrence of hiatus hernia measuring >2 cm in length, on follow-up gastroscopies, and/or trans-diaphragmatic gastric protrusion on CT-imaging. Secondary outcomes included symptom recurrence, length of hospital admission, post-operative complications and 30-day mortality. This study was approved by the local Institutional Review Board and performed in accordance with the Declaration of Helsinki.

3.2. Surgical Technique

Overweight or obese patients received preoperative low calorie and high protein diet of Optifast to reduce liver size. All patients received preoperative IV antibiotics and DVT prophylaxis with sequential calf compressors and 20mg subcutaneous enoxaparin, as well as an indwelling catheter to monitor urine output and a temporary nasogastric tube to decompress the stomach. They are placed in lithotomy and anti-Trendelenburg position. Pneumoperitoneum is created via closed Veress entry at Palmer's point. Four ports are inserted in the upper abdomen under direct vision and a Nathanson liver retractor is preferentially used. Dissection begins at the lesser curvature at the level of the pars flaccida of the gastrohepatic ligament with identification of the caudate lobe, inferior vena cava, and right crus. Hernia sac is opened and progressively dissected off the mediastinal space, with identification of the left crus. The hernia sac is completely reduced into the abdomen and excised using an ultrasonic dissection device. This is followed by circumferential mobilisation of the lower oesophagus, after careful dissection of the perioesophageal congenital adhesions within the hiatus. A nylon tape is placed around the oesophagus to aid in gentle lateral, downwards and upwards tractions. The distal oesophagus is mobilised into the mediastinum to approximately 7cm or until GOJ can be easily reduced 2-3 cm below the diaphragm without tension (Image 1). Collis gastroplasty or diaphragm relaxing incisions were not utilised. A bougie was also not utilised. Both pleurae are visualized and preserved. Diaphragmatic crura are primarily closed posteriorly with continuous size 1 Stratafix™ (Ethicon) suture and the phreno-oesophageal ligament is re-created using two interrupted 2/0 PDS sutures on both sides. Repair is reinforced with a Phasix™ non-ST mesh cut into a standardized U-shaped 7x10cm configuration, placed posteriorly and secured on the crura with 4mls of Tisseel fibrin glue (Image 2). Posterior gastropexy (close to the GOJ) to the left crus and partial 120o-150o anterior fundoplication (modified Dor) is completed with interrupted 2/0 Ethibond sutures. An intra-abdominal drain is only used when indicated. The nasogastric tube is routinely removed at the end of the procedure and patients receive regular postoperative intravenous




Image 1: A nylon tape is placed around the oesophagus to aid in gentle lateral, downwards and upwards tractions. The distal oesophagus is mobilised into the mediastinum to approximately 7cm or until GOJ can be easily reduced 2-3 cm below the diaphragm without tension.




Image 2: Repair is reinforced with a Phasix™ non-ST mesh cut into a standardized U-shaped 7x10cm configuration, placed posteriorly and secured on the crura with 4mls of Tisseel fibrin glue.

antiemetics of Ondansetron 4mg and Dexamethasone 8mg. A post-operative TTE is routinely performed the following day in HDU to exclude cardiac tamponade, and the patient is commenced on a puree diet before discharge home.

3.3. Follow-Up

Perioperative complications were defined in accordance with the modified Clavien-Dindo classification [12]. Patients were followed up at 2- and 6-weeks, and 4- and 12-months after discharge from hospital. Routine gastroscopy and CT chest for evaluation of hernia recurrence were scheduled at 12 months post-surgery, and then yearly.

4. Results

A total of 32 patients (mean age 68.8 years) underwent laparoscopic repair of large sliding hiatus or paraoesophageal hernias with modified anterior Dor fundoplication, using a Phasix™ non-ST mesh for crura reinforcement. Commonly reported preoperative symptoms included dyspnoea (62.5%), dysphagia (53.1%), epigastric pain (46.9%) and chest pain (43.7%). Mild reflux symptoms were reported in 56.3% of cases. Seven patients (21.9%) were diagnosed with a large type I sliding hiatus hernia (measuring >7 cm in length), 20 with a type III paraoesophageal hernia (62.5%) and five with a type IV paraoesophageal hernia (15.6%). No patients had a type II paraoesophageal hernia. Pre-operatively, three patients were ASA class I, ten were ASA class II, 18 were ASA class III, and one ASA class IV. Patient demographics and pre-operative characteristics are described in Table 1. All the procedures were completed laparoscopically and there were no intra-operative complications

recorded during our study. The median hospital length of stay was 3 days (range 2-7 days) and the overall post-operative complication rate was 6.3% (n=2). Both complications were recorded as grade I (one asthma attack and one basal lung atelectasis) according to the modified Clavien-Dindo scale. There were no major complications and no post-operative mortality at 30 days (Table 2). The median follow-up time was 26 months (range 12-53). No patients were lost to follow-up at the 2-year mark. No mesh-related complications were recorded and 93.8% of patients (n=30) remained symptom-free. There was only one endoscopic and radiological recurrence found in one patient at 18 months post-surgery, who was immunosuppressed on steroids for severe asthma and suffered chronic cough from bronchiectasis.

5. Discussion

In 1919, the first open hiatus hernia repair was described by Soresi[13]. Many years passed until Cuschieri described laparoscopic repair of large hiatus hernias, paving the way for considerable advances in the surgical approaches to paraoesophageal hernia repair [14]. These hernias are characterized by significant displacement of

Table 1: Characteristics of patients who underwent laparoscopic repair of large hiatus & para-oesophageal hernias with Phasix™ non-ST mesh reinforcement.

	<i>n</i>
Mean age (years)	68.8 (range 38-87)
Sex	
Male	10
Female	22
Symptoms[†]	<i>n</i>
Heartburn or reflux	18
Regurgitation or vomiting	8
Epigastric or retrosternal pain	15
Dysphagia or swallowing difficulty	17
Abdominal bloating	2
Nausea	1
Chest pain	14
Dry cough	3
Burping	2
Iron deficiency anaemia	7
Breathing difficulties or breathlessness	20
Dizziness	6
Pre-operative investigations[‡]	
Gastroscopy	32
Oesophageal manometry	1
Computed tomography (CT)	32
Barium swallow	1
Echocardiogram	12
Coronary angiogram	2
Hernia classification	
Type I	7
Type II	0
Type III	20
Type IV [§]	5
ASA class	
ASA I	3
ASA II	10
ASA III	18
ASA IV	1

[†] Note that numbers do not add up to 32 as some patients reported >1 symptom.

[‡] Note that numbers do not add up to 32 as some patients underwent >1 pre-operative investigation.

[§] Three greater omentum and two transverse colon.

Table 2: Post-operative follow up results.

	<i>n</i>
Post-operative complications	
Major	0
Minor [†]	2
Follow-up	
30-day mortality	0
Median length of stay (days)	3 (range 2-7)
Median follow-up (months)	26 (range 12-53)
Symptom-free on last follow-up	30
Endoscopic recurrence at 12 months [‡]	0
Endoscopic recurrence at 24 months [§]	1
Chest CT recurrence at 12 months	0
Chest CT recurrence at 24 months [§]	1

[†] Minor complications included lung atelectasis and asthma.

[‡] Interruption of endoscopies due to COVID-19.

[§] Hernia recurrence in the same patient.

the gastric fundus in the mediastinum, and occasionally the entire stomach and other abdominal viscera through a large defect in the phreno-oesophageal membrane. They may pose various significant and potentially life-threatening complications such as gastric volvulus, total gastric obstruction, pulmonary aspiration, and bleeding [15-16]. Surgical intervention is therefore recommended for most symptomatic patients, provided they are fit for surgery. A laparoscopic approach is now considered the gold-standard management for symptomatic paraoesophageal hernias, especially in reducing post-operative pain, length of hospital admission, and faster recovery. Despite this, the repair of giant paraoesophageal hernias remains a challenging surgical endeavour with high incidence of recurrence reported between 4%-66% in the literature.⁵⁻⁷ This remains a rather nuanced issue as most recurrences tend to be radiological in nature, with a minority resulting in symptom recurrence. Traditionally, surgical repair of paraoesophageal hernias has relied on primary suture hiatal repair which, unfortunately, has shown relatively high recurrence rates [17]. Case series evaluating modified surgical techniques such as excision of the hernia sac and simple anterior gastropexy show a variable 0%-16% recurrence rate at 24 months [18-20]. Mesh reinforcement has emerged as an alternative approach to reduce recurrence rates by reinforcing the hiatus and improving durability of the repair. While early evidence suggested a significant reduction in hernia recurrence rates, recent studies have not consistently demonstrated the superiority of mesh over primary suture hiatal repair [21]. Different types of prostheses have been used. Synthetic meshes are made of polypropylene, polyester or ePTFE, but have unfortunately demonstrated a long-term risk of oesophageal erosion or stricture [22,23]. A recent systematic review which included 35 case reports and 20 observational studies demonstrated synthetic mesh is more frequently implicated in mesh erosion, with the majority of incidences occurring within 5 years of surgery [24]. Although biological meshes (porcine small intestinal submucosal, bovine pericardium or allograft dermal matrix) do not have the same long-term risk of erosion, they have been shown to have no long-term benefit in term of hernia recurrence compared to synthetic meshes.^{8,9} Biosynthetic meshes such as Gore Bio-A®, a polyglycolide and trimethylene carbonate polymer which resorbs in 6-8 months, have been utilised with varying degrees of recurrence from 2%-18% [25,26]. The emergence of new biosynthetic meshes, such as Phasix™, may present a promising alternative in paraoesophageal hernia repair. Phasix™ mesh is a knitted monofilament mesh which fully resorbs within 12-18 months, which is longer than other comparable biosynthetic meshes available. This offers potential advantages in terms of reducing the risk of erosion associated with synthetic meshes, while avoiding the potential long-term recurrence risk observed with some other biological and biosynthetic meshes as it resorption time is slower compared to other similar meshes [27-29]. A recent 5-year outcome prospective study

published by DeMeester's group reported a 25% recurrence rate (8/32 patients) following laparoscopic paraoesophageal hernia repair with composite PhasixTM ST mesh reinforcement that comprises a hydrogel barrier on the intraabdominal side [30]. In this study, Nissen fundoplication was the treatment of choice, which doesn't secure the stomach to the diaphragm, compared to Dor fundoplication. Furthermore, they preferentially secured the mesh to the crura using absorbable tacks, which isn't without risk of vascular injury [31]. Similarly, following a median follow-up of 27 months Aiolfi et al. [32]. Reported a lower hernia recurrence rate of 8.8% (6/68) using the same mesh with a Toupet fundoplication, while Panici Tonucci and coll. published a 3.2% recurrence rate (2/62) after a median follow-up of 17 months, with Toupet fundoplication, which is almost comparable to our findings (3.1% recurrence after a median follow-up of 26 months) [32,33]. None of those three series reported any mesh-related complications. We evaluated the feasibility of PhasixTM non-ST mesh for the repair of giant paraoesophageal hernias. Similar to Gore Bio-A[®], PhasixTM non-ST mesh lacks a hydrogel barrier thus promoting adhesion formation on both sides. The aim is to purposely promote stronger connections between the hiatal repair and the posterior gastric wall, which should theoretically ensure permanent reinforcement of the repair long after the mesh has been reabsorbed, and we hypothesise that recurrence rate could be favourably reduced. Our study recorded no mesh-related complications and 93.8% of patients remained symptom-free at median 26 months follow-up, with only one demonstrating hernia recurrence (3.1%). To our knowledge, this is the first study investigating the feasibility of PhasixTM non-ST over a median >2-year follow up for large hiatal/paraesophageal hernia repairs. Although still too early to draw any conclusion, our results compare favourably to the previously mentioned studies using PhasixTM ST mesh. Only extended follow up will allow us to determine if PhasixTM non-ST can still improve long-term outcome, following laparoscopic paraoesophageal hernia repair compared to PhasixTM ST mesh. Re-do hiatal hernia repairs are technically challenging procedures with a higher risk of postoperative morbidity compared to native hiatal hernia repairs. A recent retrospective study by Liu et al. [34]. Analysed 346 patients undergoing revision hiatal surgery, of which [35] had pre-existing mesh, and noted longer operative times and higher risk of intraoperative complications such as bleeding and injury to surrounding lung, liver, and pleura (48.6 vs 22.5%) compared to the non-mesh group [34]. Of importance, this risk was only seen with non-absorbable mesh; absorbable mesh was not associated with an increased risk of complications in revisional surgery, however the retrospective review only included 9 cases with pre-existing absorbable mesh and may not have sufficient sample to draw strong conclusions [34]. Another study by Barazanchi et al. [35] had similar findings with increased risks associated with revisional hiatus surgery with pre-existing mesh [35]. However, this study only had one patient that had a pre-existing absorbable mesh. Thus, more research is needed to see if absorbable meshes have the same risk profile as non-absorbable meshes in relation to revisional hiatus hernia surgery. The PhasixTM non-ST mesh lacks a hydrogel barrier and theoretically promotes adhesions on both sides of the device, with the goal being to prevent recurrence. If a hernia were to still re-occur despite the Phasix non-ST mesh, risk of complications for a re-do should be similar to the PhasixTM ST mesh.

6. Limitations

We conducted an observational prospective cohort study at a single metropolitan hospital in Sydney, Australia. We recognise the risk of introduced bias as the population of patients undergoing paraoesophageal hernia repair is, by nature, inhomogeneous. Owing the nature of our single-centre study we can at least guarantee the consistency of the surgical technique performed throughout the entire cohort study group. We were also limited by a relatively small sample size as PhasixTM mesh only became available in Australia in 2019 and consequently the study may not have sufficient power to detect long-term mesh related complications as they tend to be uncommon.

7. Conclusion

Our single-centre early experience seems promising for medium-to-long term outcomes with PhasixTM non-ST mesh reinforcement of the oesophageal hiatus. More studies with a larger sample size and longer follow up duration are needed to further evaluate the safety profile and efficacy of this mesh in paraoesophageal hernia repairs. Due to the potentially limited number of recruited patients per specialist centre, we would recommend further collaborative prospective multi-centre randomised clinical trials involving the use of biosynthetic mesh to establish its definitive role and potential preferred option in complex hiatus hernia repair strategies, including choice of fundoplication as Nissen doesn't secure the stomach to the diaphragm, compared to Dor fundoplication.

Future research should also emphasize assessing quality-of-life outcomes related to biosynthetic mesh usage, understanding the long-term durability, and potential benefits in reducing recurrence rates beyond the observed follow-up in our study. Moreover, comparisons between different types of biosynthetic meshes, including PhasixTM ST and PhasixTM non-ST, but also other biosynthetic mesh variants, would provide valuable insights into the most optimal mesh choice for paraoesophageal hernia repair, as the debate is still opened..

References

- Kahrilas PJ. Hiatus hernias causes reflux: Fact or fiction? *Gullet*. 1993;3(Suppl):21.
- Dietrich CG, Hubner D, Heise JW. Paraesophageal hernia and iron deficiency anaemia: Mechanisms, diagnostics and therapy. *World J Gastrointest Surg*. 2021;13(3):222-230.
- Stylopoulos N, Gazelle GS, Rattner DW. Paraesophageal hernias: operation or observation? *Ann Surg*. 2002;236(4):492-500.
- Sihvo EI, Salo JA, Rasanen JV, Rantanen TK. Fatal complications of adult paraesophageal hernia: A population-based study. *J Thorac Cardiovasc Surg*. 2009;137(2):419-24.
- Ortenzi M, Balla A, Fonatana G. Factors influencing recurrence after minimally invasive treatment of hiatal hernia – a single center experience. *Laparosc Surg*. 2020;4:39.
- Hietaniemi H, Ilonen I, Jarvinen T. Health-related quality of life after laparoscopic repair of giant paraesophageal hernia: how does recurrence in CT scan compare to clinical success? *BMC Surg*. 2020;20(1):109.
- Dallemagne B, Kohnen L, Perretta S, Weerts J, Markiewicz S, Constant J. Laparoscopic repair of paraesophageal hernia. Long-term follow-up reveals good clinical outcome despite high radiological recurrence rate. *Ann Surg*. 2011;254(2):291-6.
- Petric J, Bright T, Liu DS, Wee Yun M, Watson DI. Sutured versus mesh-augmented hiatus hernia repair: a systematic review and meta-analysis of randomized controlled trials. *Ann Surg*. 2022; 275(1): e45-e51.
- Angeramo CA, Schlottmann F. Laparoscopic paraesophageal hernia repair: to mesh or not to mesh systematic review and meta-analysis. *Ann Surg*. 2022; 275(1):67-72.
- Rajkomar K, Wong CS, Gall L. Laparoscopic large hiatus hernia repair with mesh reinforcement versus suture cruroplasty alone: a systematic review and meta-analysis. *Hernia*. 2023; 27(4):849-860.
- Liu DS, Wong DJ, Goh SK. Quantifying perioperative risks for antireflux and hiatus hernia surgery: a multicenter cohort study of 4301 patients. *Ann Surg*. 2024; 279(5):796-807.
- Clavien PA, Barkun J, de Oliveira ML. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg*. 2009;250(2):187-96.
- Soresi AL. Diaphragmatic hernia: its unsuspected frequency: diagnosis and technique for radical cure. *Ann Surg*. 1919;69(3):254-70.

14. Cuschieri A. Laparoscopic antireflux surgery and repair of hiatal hernia. *World J Surg.* 1993; 17(1): 40-5.
15. Kahrilas PJ, Kim HC, Pandolfino JE. Approaches to the diagnosis and grading of hiatal hernia. *Best Pract Res Gastroenterol.* 2008;22(4): 601-616.
16. Luketich JD, Raja S, Fernando HC. Laparoscopic repair of giant paraesophageal hernia: 100 consecutive cases. *Ann Surg.* 2000; 232(4): 608-618.
17. Oelschlager BK, Pellegrini CA, Hunter J. Biologic prosthesis reduces recurrence after laparoscopic paraesophageal hernia repair. *Ann Surg.* 2006; 254(1): 36-41.
18. Van der Peet DL, Klinkenberg-Knol EC, Poza AA, Siestes C. Laparoscopic treatment of large paraesophageal hernias: both excision of the sac and gastropexy are imperative for adequate surgical treatment. *SurgEndosc.* 2000;14(11):1015-8.
19. Ponsky J, Rosen M, Fanning A, Malm J. Anterior gastropexy may reduce the recurrence rate after laparoscopic paraesophageal hernia repair. *SurgEndosc.* 2003;17(7):1036-41.
20. Daigle CR, Funch-Jensen P, Calatayud D, Rask P, Jacobsen B. Laparoscopic repair of paraesophageal hernia with anterior gastropexy: a multicenter study. *SurgEndosc.* 2015;29(7):1856-61.
21. Antoniou SA, Pointner R, Granderath FA. Hiatal hernia repair with the use of biologic meshes: a literature review. *SurgLaparoscEndoscPercutan Tech.* 2011; 28(2): 61-69.
22. Gruber-Blum S, Brand J, Keibl C. Mesh erosion and fistulation to the esophagus after incisional hernia repair with polypropylene mesh. *Hernia.* 2021; 25(5): 1169-1172.
23. Pierce RA, Spitler JA, Frisella MM. Pooled data analysis of laparoscopic vs. open ventral hernia repair: 14 years of patient data accrual. *SurgEndosc.* 2009; 23(4): 755-761.
24. Spiro C, Quarmby N, Gananadha S. Mesh-related complications in paraesophageal repair: a systematic review. *SurgEndosc.* 2020;34(10):4257-4280.
25. Gebhart A, Vu S, Armstrong C, Smith BR, Nguyen NT. Initial outcomes of laparoscopic paraesophageal hiatal hernia repair with mesh. *Am Surg.* 2013;79(10):1017-1021.
26. Silecchia G, Iossa A, Cavallaro G, Rizzello M, Longo F. Reinforcement of hiatal defect repair with absorbable mesh fixed with non-permanent devices. *Minim Invasive Ther Allied Technol.* 2014;23(5):302-308.
27. Smith JK, Puskas JD. The increasing use of biologic meshes in ventral hernia repair: benefits and limitations. *Expert Rev Med Devices.* 2012; 9(5): 511-522.
28. Novitsky YW, Harrell AG, Cristiano JA. Comparative evaluation of adhesion formation, strength of ingrowth, and textile properties of prosthetic meshes after long-term intra-abdominal implantation in a rabbit. *J Surg Res.* 2007; 140(1): 6-11.
29. ColavitaPD, Belyansky I, Walters AL. Twelve-month outcomes of abdominal wall reconstruction with a biological prosthesis: a prospective study. *Am J Surg.* 2017; 214(1).
30. McKay SC, Dunst CM, Davila-Bradley D, Reavis KM, DeMeester SR. Five-year outcomes from a prospective study on the safety and efficacy of Phasix-ST mesh use at the hiatus during paraesophageal hernia repair. *J Am Coll Surg.* 2024; 239(4):333-338.
31. Abdelmoaty WF, Dunst CN, Filicori F. Combination of surgical technique and bioresorbable mesh reinforcement of the crural repair leads to low early hernia recurrence rates with laparoscopic paraesophageal hernia repair. *J Gastrointest Surg.* 2022;24(7):1477-1481.
32. Aiolfi A, Cavalli M, Sozzi A. Medium-term safety and efficacy profile of paraesophageal hernia repair with Phasix-ST® mesh: a single-institution experience. *Hernia.* 2022;26(1):279-286.
33. Panici Tonucci T, Asti E, Sironi A, Ferrari D, Bonavina L. Safety and Efficacy of Crura Augmentation with Phasix ST Mesh for Large Hiatal Hernia: 3-Year Single-Center Experience. *J Laparoendosc Adv Surg Tech A.* 2020;30(4):369-372.
34. Liu DS, Allan Z, Wong DJ. Pre-existing hiatal mesh increases morbidity during and after revisional antireflux surgery: a retrospective multicenter study. *Surg.* 2023; 174(3): 549-557.
35. Barazanchi AWH, Chui JN, Bhimani N, Leibman S, Smith G. Impact of pre-existing mesh at the hiatus at revisional hiatal hernia surgery. *Dis Esophagus.* 2024; 37: 1-7.